



van der Nelson, H. A., Draycott, T., Siassakos, D., Yau, C. W. H., & Hatswell, A. J. (2017). Carbetocin versus oxytocin for prevention of post-partum haemorrhage at caesarean section in the United Kingdom: An economic impact analysis. *European Journal of Obstetrics and Gynecology and Reproductive Biology*, 210, 286-291. <https://doi.org/10.1016/j.ejogrb.2017.01.004>

Peer reviewed version

License (if available):  
CC BY-NC-ND

Link to published version (if available):  
[10.1016/j.ejogrb.2017.01.004](https://doi.org/10.1016/j.ejogrb.2017.01.004)

[Link to publication record in Explore Bristol Research](#)  
PDF-document

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:  
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

**Table 1: Proportionate spread of PPH events across categories of blood loss**

		Proportion of all PPHs falling in each category				Total
		No PPH (blood loss <500ml)	500-999ml	1000 – 1499ml	≥1500ml	
Holleboom 2013	Oxytocin	680	319	66	57	1122
	Carbetocin	248	165	33	16	462
Raw data from published literature						
Su 2012	Oxytocin	563	35			598
	Carbetocin	574	23			597
Model inputs, including interpolation of missing data using data from the Holleboom et al., dataset						
Su 2012	Oxytocin	94.2%	4.3%	0.9%	0.7%	100%
	Carbetocin	96.2%	2.8%	0.6%	0.4%	100%